


# C3 – Quantitative Chemistry

## Conservation of Mass

- Atoms cannot be created or destroyed during reactions.
- **Mass of reactants = mass of products.**


To show mass is conserved in a reaction:

$M_r$  on the left-side must be same as the right side.



## Reacting masses

Use conservation of mass to predict masses:



Both sides need to be equal:  
 $10\text{g} - 8\text{g} = 4\text{g}$  of oxygen

## Percentage Mass

- Percentage mass of an element in a compound

$$\frac{\text{Mass of the element in compound}}{\text{Total mass of compound}} \times 100$$

**Example Question:**

**Find the percentage mass of oxygen in magnesium oxide (MgO).**

$A_r$  of magnesium = 24       $A_r$  of oxygen = 16

$M_r$  of MgO =  $24 + 16 = 40$

$$\begin{array}{rcl} \% \text{ mass} = \frac{A_r}{M_r} = \frac{16}{40} = 0.4 \times 100 = 40\% \end{array}$$

X 100 to make a %      40% of the mass of MgO is oxygen


## Mass Changes

- Mass is always conserved in a reaction.
- Sometimes it may seem like the mass has increased/decreased.
- If a **reactant** is a gas – mass may **increase**.




Oxygen is in the air before it combines with magnesium – you cannot find the mass of oxygen on the balance.

It will look like the mass has increased when it is re-weighed at the end.



- If a **product** is a gas and the gas is able to escape the system – mass will **decrease**.




It will look like the mass has decreased as some of the atoms have been given off as gas and have escaped – so cannot be re-weighed.

## Atomic mass ( $A_r$ ) and Relative Formula Mass ( $M_r$ )

- Atomic mass ( $A_r$ ) is the mass number – ie the mass of one atom
- Relative formula mass ( $M_r$ ) = all the **relative atomic masses ( $A_r$ )** of the atoms in a compound or molecule added up.

**Example**



## The Mole (HT only)

- **Avogadro constant –  $6.02 \times 10^{23}$**
- One mole contains  $6.02 \times 10^{23}$  atoms or molecules
- The mass, in g, of one mole is the  $A_r$  (if an element) or  $M_r$  if a compound or molecular element

$$\text{Mass} = M_r \times \text{moles}$$

Iron has a  $A_r$  of 56, so 1 mole of iron is 56 g and contains  $6.02 \times 10^{23}$  atoms of iron

Ammonia ( $\text{NH}_3$ ) has an  $M_r$  of 17, so 1 mole of ammonia has a mass of 17g. and contains  $6.02 \times 10^{23}$  molecules of ammonia

### C3 – Quantitative Chemistry

1. What is meant by conservation of mass?
2. Mass of reactants = ?
3. The  $M_r$  of the left side of an equation must be the same as..

1. How do you calculate the percentage mass of an element in a compound?
2. What do you do to convert a decimal into a percentage?

1. Should mass change in a reaction?
2. If a reactant is a gas, what will happen to the mass?
3. Why will it appear this has happened?

1. What does  $M_r$  stand for?
2. What is the relative formula mass?
3. Where can you find the relative atomic mass ( $A_r$ ) of an element?

1. How many atoms are in one mole?
2. How do we know what the mass of one mole of an element is?
3. How do we know the mass of one mole of a compound?

4. If a product is a gas, what will happen to the mass?
5. Why will it appear this has happened?

# C3 – Quantitative Chemistry

## Concentrations of Solutions

- Concentration = mass of dissolved substance in specific volume (eg dm<sup>3</sup>)
- More substance dissolved = more concentrated solution

$$\text{Concentration} = \frac{\text{mass}}{\text{volume}}$$

(g/dm<sup>3</sup>)      (g)      (dm<sup>3</sup>)

Can be rearranged to find mass dissolved:

$$\text{mass} = \text{concentration} \times \text{volume}$$

(g)      (g/dm<sup>3</sup>)      (dm<sup>3</sup>)

$$1000\text{cm}^3 = 1\text{dm}^3$$

$$\text{cm}^3 \rightarrow \text{dm}^3 = \text{divide by } 1000.$$

## Calculating mass in a given volume

If you have a known volume of a solution of known concentration then you can calculate the mass of dissolved solid.

E.g Calculate the mass of dissolved solid in 25cm<sup>3</sup> of a 96g/dm<sup>3</sup> solution

96g/dm<sup>3</sup> means 96g in every 1000cm

Do the same to the other side (÷40)

↓  
2.4g

25cm<sup>3</sup>

How do we get from 1000 to 25? (÷40)

## Moles and Equations (HT only)

- You can use moles to help you write balanced symbol equations.

### Example Question

18.4g of Sodium reacted with 6.4g of oxygen to give 24.8g sodium oxide. Use the masses to write the balanced equation.

Step	Example
Write the equation for the reaction (unbalanced)	Na + O <sub>2</sub> → Na <sub>2</sub> O
write down the mass or % given in the question	18.4 + 6.4 → 24.8
Write the mass of one mole of each element or compound	23      32      62 (e.g 18.4 ÷ 23)
Divide the mass given in question by the mass of one mole	0.8      0.2      0.4
Turn the answers into whole number simple ratio	8      2      4 (cancel down) 4      1      2
Put the numbers into the equation	4Na + O <sub>2</sub> → 2Na <sub>2</sub> O

## Calculating reacting masses (HT)

### Example Question

Calculate the mass of calcium needed to make 11.2g Calcium oxide

Step	Calculation
Write the balanced equation	2Ca + O <sub>2</sub> → 2CaO
Write the masses of each substance	80 + 32 → 112
Write down the given mass in the question.	11.2
Work out the 'scale' factor (ie what did you have to do to the original number to get to the desired mass	÷ 10
Do the same to the other side	8g

## Limiting Reactants (HT only)

- If one reactant runs out before the other, then the reaction will stop.
- The reactant that runs out first in a reaction is known as the limiting reactant.

## C3 – Quantitative Chemistry

1. What does concentration mean?
2. How can you make a solution more concentrated?
3. State the equation to calculate concentration in  $\text{g/dm}^3$ .
4. What is the unit for volume?
5. How many  $\text{cm}^3$  are in a  $\text{dm}^3$ ?

### Calculating mass in a given volume

1. What does  $36.5\text{g/dm}^3$  mean?
2. Calculate the mass of dissolved solid in  $25\text{ cm}^3$  of a  $36.5\text{g/dm}^3$  solution

36.5

1000

25  $\text{cm}^3$

Do the same to the other side  
( $\div 40$ )

g

How do we get from 1000 to 25?  
( $\div 40$ )

### Moles and Equations (HT only)

12g of magnesium (Mg) reacted with 8g of oxygen ( $\text{O}_2$ ) to produce 20g magnesium oxide (MgO). Use the masses to write a balanced equation

Step	Example
Write the equation for the reaction (unbalanced)	
write down the mass or % <u>given in the question</u>	
Write the mass of one mole of each element or compound	
Divide the mass given in question by the mass of one mole	
Turn the answers into whole number simple ratio	
Put the numbers into the equation	

1. What is a limiting reactant?
2. Complete the calculation:  
Calculate the mass of calcium needed to make 224g of calcium oxide

Step	Calculation
Write the balanced equation	$2\text{Ca} + \text{O}_2 \rightarrow 2\text{CaO}$
Write the masses of each substance	
Write down the given mass in the question.	
Work out the 'scale' factor (ie what did you have to do to the original number to get to the desired mass)	
Do the same to the other side	